

A New Possible Way to Explain the DAMA Results

J.Va'vra

SLAC, Stanford University, CA94309, U.S.A.
e-mail: jjv@slac.stanford.edu

Received January 10, 2014

ABSTRACT

At present there is an effort to reconcile the results of the DAMA experiment with those from other Dark Matter experiments such as CoGeNT, CRESST, CDMS, and all LXe experiments. The author suggests a new model describing the Dark Matter signal as the result of collisions of very light (1-to-few GeV/c^2) WIMPs with hydrogen, and compares it with currently accepted models of collisions with heavy nuclei (Na, Ge or Xe). The hydrogen target would come from H-contamination of NaI(Tl), Ge and CaWO_4 crystals. Initial tuning indicates that one can explain the modulation amplitude of DAMA and CoGeNT with this model, assuming a WIMP-proton cross section between 10^{-33} and 10^{-32} cm^2 . This paper should be considered to be a new idea which will need substantial new experimental input from all involved experiments.

Key words. DAMA experiment, Dark Matter search

1. Introduction

The DAMA experiment observes a small, but clear, oscillatory signal. The signal is a peculiar modulation of the counting rate of 2-6 keV single-hit events (i.e. those in which only one of the many detectors in the setup fires) satisfying all the many requirements of the Dark Matter annual modulation signature. The observed yearly modulation is in phase with the Earth's motion around the Sun. DAMA is the first experiment to observe a clear oscillatory signal [1] and [2], as shown on Fig. 1.

The CoGeNT experiment (see Fig. 2) also reports evidence of oscillation, and the CRESST-II and CDMS experiments also report hints of a signal. However, there is no hint of a signal from any LXe Dark Matter Search experiments, such as Xenon-10, Xenon-100 or LUX.

Figure 3, which was shown at the 2013 Snowmass meeting, presents one of the possible model-dependent comparisons to the results of dark matter experiments. The important point for this paper is that there is great opportunity to explore the cross section region of $10^{-30} - 10^{-44} \text{ cm}^2$ if the mass is indeed low.




Fig. 1. DAMA data: residual rate of the single-hit scintillation events, measured by DAMA/LIBRA over six annual cycles in the (2-6) keV energy interval as a function of time [2]. The modulation amplitude is ~ 0.01 counts/day/kg/keV.




Fig. 2. CoGeNT data: counts per 30-day interval from the 0.5-to-few keVee energy window [3]. The modulation amplitude is ~ 0.3 counts/day/kg/keV.




Fig. 3. The model-dependent status of cross section limits for the detection of the WIMP mass from various experiments (Snowmass, 2013). The graph also shows the low limit, below which experiments will have to deal with neutrino background. One can see clearly, that a WIMP mass below $\sim 5 \text{ GeV}/c^2$ would not be detected by most of the present experiments.




Fig. 4. As for Fig. 3, but the figure also shows the future reach of the CDMS experiment at low mass (presented by CDMS at Snowmass, 2013).¹

Figure 4 shows the CDMS future low mass limit reach¹ for various new detector configurations and using new improved electronics. Clearly, CDMS aims to explore the low mass range, and so the idea presented in this paper will be tested soon.

2. A new idea to explain the DAMA data

The author discusses a new idea, which explains the DAMA signal as being due to the scattering of a light Dark matter particle (1-to-few GeV/c^2) on a light nucleus; specifically on hydrogen present as a small OH or H_2O contamination in the NaI(Tl) crystals.

Table 1 shows the maximum calculated kinematical limit in terms of the nuclear recoil (keVnr) and the electron-equivalent recoil (keVee).² We point out that DAMA quotes the threshold in keVee units, so that one has to convert keVnr units to keVee units. One can easily calculate the keVnr-scale using two-body kinematics; to convert the keVnr-energy-scale to the keVee-scale, one needs to know the quenching factor (QF) for a given nucleus, i.e., $E_{keVee} = \text{QF} * E_{keVnr}$. The quenching factor depends on the recoil energy. In case of NaI(Tl), for sodium recoil, I assume $\text{QF} \sim 0.25$ with the understanding that the value could be lower.³ The proton quenching factor in NaI(Tl) has not been measured to our knowledge.⁴ Generally it is expected that the proton QF is considerably higher than for Na or I recoils, but not quite as high as 1.0. In the absence of any measurement, we follow the semi-empirical calculation of V. Tretyak [6], which is

¹ B. Cabrera, private communication; figure presented at the Snowmass meeting, 2013 on behalf of the CDMS group.

² It should be noted that the literature indicates quite a scatter in values of the Galactic escape velocity: (a) $450 \text{ km/sec} < v_{esc} < 650 \text{ km/sec}$ [8], (b) $489 \text{ km/sec} < v_{esc} < 730 \text{ km/sec}$ [9], and (c) $498 \text{ km/sec} < v_{esc} < 608 \text{ km/sec}$ [10].

³ Some experiments measured the quenching factor QF for sodium in NaI(Tl) close to 0.25-0.3, see for example [4] and [6]. However, according to private communication with J. Collar, some more recent measurements indicate values close to 0.1-0.15 for a recoil energy of $\sim 10 \text{ keVnr}$ [5]; recently this has been measured independently at Duke University, with a similar result (soon to be published). However, as discussed in the paper, the QF factor depends on exact experimental conditions, i.e., it is not a universal constant [6].

⁴ Confirmed by private communication with J. Collar, D. McKinsey and V.I. Tretyak.




Fig. 5. The semi-empirical calculation⁵ of the quenching factor for proton recoils in the NaI(Tl) crystal, for two values of the Birks constant. DAMA determined a low value of $kB = 1.25 \text{ mg}/(\text{cm}^2 \text{ MeV})$ using alpha particles measured in their experiment; this in turn gives a higher QF value for very small proton recoil energies.⁵

based on only one parameter, kB , the so-called Birks factor. The value of the kB factor depends [6] on many parameters, such as the detector material itself with all its impurities and dopants, the conditions of the experiment such as temperature, the signal integration time, etc. Ideally, the QF value should be measured in the same experiment, and under exactly the same conditions as used for data taking. The kB factor could be determined by fitting experimental data obtained using one type of particle, for example α particles from an external source, and then using it to predict the QF value for another type of particle, for example the proton, even in a different energy region. Figure 5 shows V. Tretyak's QF calculation⁵ for the proton QF for two values of the Birks constant, based on two extreme values obtained from different experiments; for kB constant at extreme values between 1.25 and 6.5 $\text{mg}/(\text{cm}^2 \text{ MeV})$, the proton QF is between 0.95 and 0.8 at $\sim 0 \text{ keV}$; for the DAMA case, where kB was determined to be 1.25 $\text{mg}/(\text{cm}^2 \text{ MeV})$ using the α particle calibration, the proton QF is ~ 0.95 for very small proton recoil energies. I assume a value of $\text{QF} \sim 0.95$ for the calculations of Table 1. The conclusions would not change, however, if one would take $\text{QF} \sim 0.8$ instead.

The DAMA experiment detection software threshold was 2 keVee during most of data taking [7], but it was recently lowered to 1 keVee [11] (while the hardware threshold was held at 1 photoelectron all along). Notice in Table 1 that the maximum electron-equivalent recoil energy for Na-nucleus is below the new DAMA detection threshold unless the WIMP mass is above $\sim 4 \text{ GeV}/c^2$. On the other hand, it is possible to detect a larger WIMP mass than $\sim 1 \text{ GeV}/c^2$ for scattering off the H-nucleus even with the old DAMA threshold of 2 keVee, assuming that the proton QF is ~ 0.95 .

DAMA NaI(Tl) crystals are made of Ultra-Low-Background (ULB) materials to guarantee low radioactive background. While the DAMA experiment does quote very low radio-purity levels [7], it does not quote the H, OH and H_2O contamination at all. Generally, the H-contamination is not quoted by crystal producers, since spectroscopic properties of NaI(Tl) crystals are not affected by it up to a level of $\sim 10 \text{ ppm}$. The OH-molecule represents a likely source of contamination of the primary NaI salt; its removal requires prolonged pumping at elevated temperature [12]. Figure 6 shows typical OH-contamination levels in NaI(Tl)

⁵ Private communication with V.I. Tretyak. The same calculation was used in [6].

Table 1. Maximum calculated nuclear recoil energy E_{keVnr} and maximum electron-equivalent recoil energy E_{keVee} as a function of WIMP mass, for two targets, hydrogen and sodium. It is assumed that the WIMP velocity relative to Earth is between 0 and the Galactic escape velocity of ~ 650 km/sec, all modulated by the Earth's velocity of 230 ± 30 km/sec in the Galactic coordinate system. In this table, for sodium recoil I used $QF = 0.25$ [4], and for hydrogen $QF = 0.95$.⁵

WIMP mass [GeV/c ²]	Nucleus	E_{keVnr} [keV]	E_{keVee} [keV]
0.5	H	1.91	1.81
1.0	H	4.30	4.08
1.5	H	6.20	5.89
2.0	H	7.65	7.26
2.5	H	8.78	8.34
3.0	H	9.68	9.20
0.5	Na	0.19	0.05
1.0	Na	0.73	0.18
1.5	Na	1.57	0.39
2.0	Na	2.67	0.67
2.5	Na	4.00	0.60
3.0	Na	5.53	1.38
4.0	Na	9.07	2.26

crystals⁶; the FTIR spectrum shows the OH-contamination for a crystal located in a protective enclosure (solid curve) just after its production, and the H₂O contamination when it was exposed to air (dashed curve). The exact level of the OH or H₂O contamination in NaI(Tl) crystals is generally proprietary information, as told by Saint-Gobain Co. However, we learned from the Hilger Co. that an OH-contamination at a level of 1-3 ppm is very likely. Generally, these companies keep the air humidity at less than $\sim 2\%$ when machining a crystal and assembling an NaI(Tl) detector, although again this is proprietary knowledge. But, knowing how hard it is to keep humidity low in gases without appropriate filtration or without pumping while baking, it would seem to us that water contamination at a level of a few ppm is easily possible. Once in the detector, each DAMA NaI(Tl) crystal is encapsulated in a copper housing and sealed for all its life. In addition, the detectors are continuously maintained in high purity nitrogen atmosphere. Therefore, if there is any H-contamination, it is probable that it would have occurred during previous manufacturing steps.

While the possible presence of H/OH-contamination in the Ultra-Low-Background (ULB) NaI(Tl) DAMA detectors remains an open question, for our calculation we will assume, in an arbitrary way, that the H-contamination is at a level of 1 ppm. Once a better number is known, one can easily scale our results.

3. Other Dark Matter search experiments

If the proposed H-contamination mechanism works for the DAMA experiment, it must also work for CDMSlite [19], CoGeNT [3], and CRESST-II [20]. The peak modulation amplitude for DAMA is ~ 300 events/30 days, for CoGeNT it is ~ 10 events/30 days; CDMSlite measures ~ 2.9 counts/keV/kg-day (no clear oscillation measured yet).

Both the CDMSlite and CoGeNT experiments use ultra-pure Ge-crystals [16]. Hydrogen is the only gas which has been successfully used for high-purity Ge-crystal growth. All commercial detector grade germanium is grown exclusively in hydro-

⁶ N. Shiran, A. Gektin, Institute of Scintillating Materials, Kharkov, Ukraine, private communication, 2014.




Fig. 6. FTIR spectrum of NaI(Tl) crystals showing the OH-contamination in a crystal located in a protective enclosure (solid curve) and the H₂O contamination when exposed to air (dashed curve).⁶

gen. Therefore, even the ultra-pure Ge has a H-contamination at a level of $\sim 2 \times 10^{15}$ atoms/cm³, or ~ 50 ppb [16], and according to P. Brink⁷ this is a reasonable estimate to assume. However, we stress that no real measurement was made on the actual crystals from these two experiments, and so we encourage that this is done. For our calculation we assume a level of ~ 50 ppb.

The CRESST-II experiment uses CaWO₄ crystals of total weight ~ 10 kg. These have been prepared by the Czochralski growth method using the solid state reaction from CaCO₃ and WO₃ from Alfa-Aesar materials with purity levels of 99.999 % and 99.998 % respectively [17]. The paper does not mention the H-contamination level, but the purity level of the major ingredients used to form the crystal is not very high, from our perspective. Reference [18] describes another possible method to prepare CaWO₄ crystals, and they did measure a significant level of OH contamination using the FTIR method. However, we were not able to obtain information on the H-contamination for actual CRESST-II crystals, even after contacting members of that collaboration. Therefore, for our calculation we assume, in an arbitrary way, an H-contamination at a level of 100 ppb. Again, once a better value is known, our results can be easily scaled.

4. Rate calculation

A low mass WIMP of 1-to-few GeV/c² represents a real experimental challenge at present for most experiments, as it requires a low mass hydrogen target in order to be detectable. So far, existing experiments provide the hydrogen target only as a small "unwanted" contamination, which limits the measured rate. Alkali halide inorganic crystals, such as NaI(Tl) or CsI(Tl), may have a unique advantage over other methods in detecting a very-low-mass WIMP, if they have sufficiently high OH-contamination.

We calculate the rate as follows:

$$Rate = \sigma_{WIMP} \cdot F_{WIMP} \cdot N_{target} \cdot V_{target} \cdot Time \cdot Det_{ef} \quad (1)$$

where Det_{ef} is the detection efficiency, σ_{WIMP} is in [cm²], F_{WIMP} is flux in [cm⁻²sec⁻¹], N_{target} is number of target particles in [cm⁻³], V_{target} is target volume in [cm³] and Time is in

⁷ P.Brink, CDMS collaboration, private communication, 2014.




Fig. 7. This paper's prediction for the modulation peak rate in several experiments as a function of cross-section, assuming the H-contamination as indicated, WIMP-proton scattering hypothesis, Dark matter density $\sim 0.3 \text{ GeV/cm}^3$ in our nearby Universe [21]. The modulation amplitude peak was calculated from a differential rate based on the Earth's velocity variation of $230 \pm 30 \text{ km/sec}$ in the Galactic coordinate system. The vertical lines indicate a cross-section corresponding to DAMA (dashed line) and to CoGeNT (solid line) measured peak amplitude rates.

Table 2. Input numbers used in the calculations for Fig. 7 and Table 3.

Experiment	weight [kg]	H-contamination	Detection efficiency	Energy range [keVee]
DAMA	250	1 ppm	0.6	1-6
CoGeNT	0.444	50 ppb	1.0	0.5-3
CDMSlite	0.6	50 ppb	1.0	0.5-3
CRESST-II	10	100 ppb	1.0	0.5-3

Table 3. Calculated number of events expected over 30 days in the peak of the modulation, for DAMA, CoGeNT, CDMSlite and CRESST-II experiments using the H-contamination model.

Cross section [cm^2]	DAMA	CoGeNT	CDMSlite	CRESST-II
10^{-32}	794	3.25	4.69	146.6
10^{-33}	79.4	0.33	0.47	14.7
10^{-34}	7.9	0.03	0.047	1.47

[sec]. The WIMP-proton scattering cross section is not known at present. We will treat it as an independent variable. Table 2 lists the input numbers used in the calculation.

Figure 7 shows our prediction for the modulation peak rate in several experiments as a function of cross section, assuming the H-contamination indicated in the figure, as a function of the WIMP-proton cross section. In order to obtain the DAMA modulation peak of ~ 300 events/30 day period a cross section slightly lower than $\sim 10^{-32} \text{ cm}^2$ is required. However, to explain the CoGeNT count rate, a slightly higher value of the cross section is required. It is therefore crucial that both experiments determine the H-contamination in their crystals.

Table 3 shows the number of events expected per 30 days in the modulation peak amplitude for four experiments.

In order to make a quantitative comparison to the proposed model, each of the four experiments should perform an H-contamination analysis of the crystals used in that experiment; DAMA should also find out if there is a correlation between rate/crystal and the H-contamination in each crystal (if indeed there is a variation in rate/crystal).

The calculation in Figure 7 is made simply to get a feel for the problem, and to provide motivation to measure the hydrogen contamination in the existing crystals. The exact tuning of the




Fig. 8. Upper limits of WIMP-nucleon cross-sections as a function of the WIMP mass, plotted for Tevatron [22] and LHC [23], CMS [24]. There is a large range of possible values of cross-section limits for $\sim 1 \text{ GeV}/c^2$ WIMP.

dark matter particle mass between 1 and a few GeV/c^2 should be done with a proper fit, which includes a realistic simulation of the dark matter velocity distribution in the Galaxy, the cross section, and the H-contamination. An example of such a calculation is found in Ref. [13], and this disagreed with our hypothesis of hydrogen contamination, as presented in our original paper [14]. However, these authors assumed a Maxwellian distribution of dark matter velocities in our Galaxy. We would argue that the dark matter velocity distribution may have two components in reality, one component representing the Maxwellian distribution, and the second component representing a direct external non-Maxwellian flow distribution, as all galaxies may be interconnected by streams of dark matter with velocities above the Galaxy escape velocity. The direct flow component may enhance the reach to a lower WIMP mass range than that which is shown in Table 1. This problem should clearly be studied in more detail.

Reference [13] also pointed out a possible conflict with a model of overheating of the Earth's atmosphere if low mass dark matter particles have too large a scattering cross section [15]. However, from Ref. [15] we conclude that we are still below their quoted overheating limits, for the proposed dark matter masses near $\sim 1 \text{ GeV}/c^2$ and a cross section below $\sim 10^{-32} \text{ cm}^2$. However, if the local dark matter density increased by a factor of ~ 10 , or if the WIMP-proton scattering cross section is larger, the overheating described in Ref. [15], might occur.

The LXe and LAr experiments do not have hydrogen contamination, and therefore any DM signal detected by these methods would directly contradict the proposed model.

5. Comparison with LHC experiments

It is interesting to compare our prediction for the DAMA WIMP-nucleon cross section range with upper limits obtained by Tevatron and LHC experiments, where the WIMP is measured in " $\overline{WIMP} + WIMP + \text{jets}$ " events [22]. This is shown in Figure 8. It is clear that the accelerator-based cross section upper limits are lower than values obtained from the H-contamination model. However the limits of collider experiments are derived with an effective operator approach, and direct comparison with our model has some unknown model dependency.

6. A new proposal for future experiments

One could attempt to detect diatomic molecular vibration, excited by gentle WIMP-proton scattering, and hence further reduce the threshold on the WIMP mass [25]. To excite such vibrations, a very small energy deposit at a level of 1.8-4.3 eV is needed. Some fraction of de-excitations goes in the visible range, so it should be possible to detect it with the Bialkali photocathode; an IR-responsive photocathode would provide an even-more-efficient method. To do this, one could use, for example, so called "wet" fused silica, which has ~ 1000 ppm of OH-contamination by design. Or, one could use simply pure water, and "tune" for the OH-absorption lines. This avenue is presently not pursued in underground experiments because of high single photo-electron noise. It could be considered in accelerator-triggered beam dump experiments searching for dark matter production. Pilot studies of this idea are in progress.

7. Conclusion

This paper suggests that the measured oscillation in the DAMA experiment may be caused by scattering of a light mass WIMP (mass between ~ 1 and a few GeV/c^2) on a hydrogen nucleus, present as a consequence of H-contamination in the NaI(Tl) crystals.

For our choice of parameters for the H-contamination model, and taking into account the DAMA modulation peak rate of ~ 0.01 cpd/kg/keV, the observed behavior can be explained with a WIMP-proton scattering cross section between 10^{-33} cm^2 and 10^{-32} cm^2 (see Fig. 7).

As the H-contamination may vary from crystal-to-crystal in the DAMA experiment. Crystal-dependent rates may be observed. This should be investigated.

We suggest that the DAMA, CDMS, CoGeNT, and CRESST-II collaborations perform an FTIR analysis on their crystals in order to measure the level of H-contamination. This will prove or disprove our claim that a single WIMP-proton scattering cross section can explain all of the data.

LXe or LAr experiments do not have hydrogen contamination, and therefore any DM signal detected by these methods would directly contradict the model proposed in this paper.

While it is somewhat puzzling that the idea presented in this paper requires a very small dark matter mass with a cross section many orders of magnitude larger than that for a particle on the mass scale of the W and Z bosons, there is nothing *a priori* which, to our knowledge, prevents this from occurring, and so we suggest that the idea be tested very carefully.

8. Acknowledgements

I also would like to thank CDMS people for discussing issues about the dark matter detection, especially Prof. B. Cabrera, S. Yellin, P. Brink, R. Partridge, G. Godfrey, and M. Kelsey. I also thank Farinaldo Da Silva Queiroz and Chris Kelso for discussions regarding reconciliation of ideas of this paper and their publication [13]. Comments of Prof. C. Galbiati encouraging to investigate this problem carefully are also appreciated. I would like thank Prof. E. Nappi for showing interest in the idea, sharing it with Prof. C. Brogini, and asking very good questions about early version of the paper. I would like to thanks Prof. E.V. Kolb for providing several references, which allowed me to enter the up-to-date status of the collider WIMP searches.

References

- [1] R. Bernabei *et al.*, Eur. Phys. J. C 23, 61 (2002).
- [2] R. Bernabei *et al.*, Eur. Phys. J. C 73, 2648 (2013).
- [3] C.E. Aalseth *et al.*, Phys. Rev. D 88 1, 012002 (2013).
- [4] G. Gerbier *et al.*, Astroparticle Physics II 287 (1999).
- [5] J. Collar, Phys. Rev. C 88, 035806 (2013), and Phys. Rev. Letters 110, 211101 (2013).
- [6] V.I. Tretyak, Astroparticle Physics 33, 4053 (2010).
- [7] R. Bernabei *et al.*, Nucl. Instr. & Meth. A 592, 297 (2008).
- [8] P.J.T. Leonard and S. Tremaine, ApJ, 547, 228 (1990).
- [9] C.S. Kochanek, ApJ, 353, 486 (1990).
- [10] M.C. Smith *et al.*, arXiv:astro-ph/0611671v2, 2012.
- [11] R. Bernabei *et al.*, JINST 7, 03009 (2012).
- [12] K.A. Kudin *et al.*, Functional Materials 18, No.2 (2011).
- [13] S. Profumo and F. Queiroz, JCAP 1405, 038 (2014).
- [14] J.Va'vra, A New Possible Way to Explain DAMA Results, arXiv:1401.0698v1, 2014.
- [15] G.D. Mack *et al.*, Phys. Rev. Lett. 112, 041302 (2014).
- [16] W.L. Hansen *et al.*, IEEE Trans. on Nucl. Sci., Vol. NS-29, No.1, 1982.
- [17] A. Erb and J-C. Lanfranchi, The Royal Society of Chemistry, CrystEngComm, 15, 2301 (2013).
- [18] Q. Li *et al.*, Hindawi Publ. Co., Journal of Chemistry, Article ID 952954 (2013).
- [19] R. Agnese *et al.*, Phys. Rev. Lett. 112, 041302 (2014).
- [20] G. Angloher *et al.*, Eur. Phys. J. C 72, 1971 (2012).
- [21] R. Catena and P. Ullio, JCAP 1205, 005 (2012).
- [22] M. Beltran *et al.*, JHEP 1009, 037 (2010).
- [23] ATLAS collaboration, <https://atlas.web.cern.ch/Atlas/GROUPS/PHYSICS/CONFNOTES/ATLAS-CONF-2011-096>.
- [24] CMS Collaboration, CMS PAS EXO-12-048, 2013.
- [25] J.Va'vra, Molecular excitations: a new way to detect Dark matter, arXiv:1402.0466v1, 2014.